

CHROMOSOME COUNTS IN THE VARIETIES OF
SOLANUM TUBEROSUM AND ALLIED
WILD SPECIES

HUGH BURNICE SMITH

Department of Biology, Illinois Wesleyan University, Bloomington, Illinois

PART II

OF A DISSERTATION

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENT FOR
THE DEGREE OF DOCTOR OF PHILOSOPHY IN THE
UNIVERSITY OF MICHIGAN

Reprinted from GENETICS 12: 84-92, January, 1927

GENETICS

A Periodical Record of Investigations Bearing on
Heredity and Variation

EDITORIAL BOARD

DONALD F. JONES, MANAGING EDITOR
Connecticut Agricultural Experiment Station

WILLIAM E. CASTLE
Harvard University

ROLLINS A. EMERSON
Cornell University

EDWIN G. CONKLIN
Princeton University

HERBERT S. JENNINGS
The Johns Hopkins University

CHARLES B. DAVENPORT
Carnegie Institution of Washington

THOMAS H. MORGAN
Columbia University

BRADLEY M. DAVIS
University of Michigan

RAYMOND PEARL
The Johns Hopkins University

EDWARD M. EAST
Harvard University

GEORGE H. SHULL
Princeton University

Genetics is a bimonthly journal issued in annual volumes of about 600 pages each.

Subscription, \$6.00 a year for complete volumes (January-November). Parts of volumes are to be had only at the single number rate. Foreign postage, 50 cents additional. Single copies, \$1.25 each, postpaid.

Volume I, Nos. 1 and 2 can no longer be supplied by the publishers. A limited number of sets of back volumes are available, with the exception of these two numbers.

Back volumes 2-10 may be had (in continuous series only) at \$7.00 each, plus postage. The Business Manager will supply information on request as to odd volumes and numbers available.

Correspondence concerning editorial matters should be addressed to Dr. DONALD F. JONES, P. O. Drawer 1106, New Haven, Conn.

Business Correspondence, including change of address, directions concerning reprints, and exchanges, should be addressed to GENETICS, Menasha, Wisconsin, U. S. A., or BROOKLYN BOTANIC GARDEN, 1000 Washington Ave., Brooklyn N. Y., U.S.A.

Remittances should be made payable to GENETICS.

Entered as second-class matter, August 31, 1922, at the post-office at Menasha, Wisconsin, under Act of March 3, 1879.

CHROMOSOME COUNTS IN THE VARIETIES OF
SOLANUM TUBEROSUM AND ALLIED
WILD SPECIES

HUGH BURNICE SMITH

Department of Biology, Illinois Wesleyan University, Bloomington, Illinois

Reprinted from GENETICS 12: 84-92, January, 1927



CHROMOSOME COUNTS IN THE VARIETIES OF *SOLANUM TUBEROSUM* AND ALLIED WILD SPECIES¹

HUGH B. SMITH

Illinois Wesleyan University, Bloomington, Illinois

Received June 21, 1926

As far as has been determined, the species in the *Solanaceae* have a basic haploid chromosome number of 12. BÖNICKE (1911) has shown that the haploid chromosome number in both *Datura Stramonium* L. and *Datura Tatula* L. is 12. BLAKESLEE (1921) reports tetraploid mutations in *Datura Stramonium* with the haploid chromosome number 24. BLAKESLEE and BELLING (1924) produced experimentally, by means of cold, tetraploid plants of *Datura Stramonium*. WINKLER (1910) reports that the chromosome number is 36 in the haploid cells of *Solanum nigrum* L. and 12 in the haploid cells of *Solanum lycopersicum* L. WINKLER (1916) produced tetraploids artificially in these two *Solanum* species from dehydrated callus tissue and as graft hybrids. From callus tissue dehydrated with chloral hydrate, adventitious buds arose whose cells contained the 4X chromosome number. In his grafting experiments, a tetraploid sprout was produced at the union of a wedge-shaped piece of nightshade scion with tomato stock. After union had taken place, a cut was made at the level of the union so that the tomato tissue was in the middle of the cut surface and the nightshade tissue was on both sides. All axial buds were removed to allow adventitious buds to grow on the cut surface. Periclinal and sectorial chimeras appeared and in one case a periclinal chimera proved to be a tetraploid branch. WINKLER produced a true tetraploid tomato from this branch by cutting the stem across and forcing adventitious buds from callus tissue. The cells of the tetraploid tomato branch contained 48 chromosomes. In somewhat the same way WINKLER obtained a giant nightshade plant whose cells contained 4X or 144 chromosomes. In connection with the work of BLAKESLEE and of WINKLER in the experimental production of tetraploid plants in the *Solanaceae*, the work of NĚMEC (1899) with potatoes is of interest. NĚMEC studied the regeneration layer which forms over cut potato tubers and reports that the chromosome number found in the cells of such tissue is surely more than 70.

¹ Papers from the Department of Biology of the MAINE AGRICULTURAL EXPERIMENT STATION, No. 177, and from the Department of Botany of the UNIVERSITY of MICHIGAN, No. 244.

The relative ease with which tetraploidy has been experimentally produced in Solanaceae would suggest that the customary vegetative reproduction of potatoes might easily, of itself, afford an opportunity for the occurrence of tetraploid mutations. By the term tetraploidy, as used in this paper, is meant the doubling of the primitive chromosome group; just how this takes place is not known at the present writing. In view of the fact that, as this paper shows, tetraploidy does occur in some of the cultivated potatoes, it is planned to attempt the artificial production of potato tetraploids from the available diploid types. Such work may give rise to the production of commercial strains of economic value. Tetraploid forms of plants are usually *gigas* forms, and *gigas* forms may possibly produce a greater yield or different quality of tubers than parental types.

It does not appear unlikely that a cytological examination of the cultivated potatoes and their wild allies may determine whether or not tetraploidy has been one of the factors in the evolution of the potato. This is the problem which is dealt with in a preliminary way in this paper. In potatoes the question of pollen sterility is a very important one. It is not impossible that one of the manifestations of sterility may be that some pollen mother cells, failing to undergo the reduction divisions produce non-functional pollen grains, whereas other microspores produced in the same anther by the normal reduction division are functional. It may also be that in certain potato varieties and species-hybrids the chromosomes are irregularly distributed to the two poles during meiosis. Although to date no study has been made of somatic material, such a study is planned for the coming season and thus evidence may be obtained for the solution of the problems suggested. It would be decidedly interesting to find out what plant breeding advances might come about through the repeated intercrossing of tetraploid plants with diploid, with the end in view of securing types with one or more reduplicated chromosomes, thus applying to a vegetatively propagated economic plant some of the results obtained by BLAKESLEE and BELLING with *Datura*.

The material from which this cytological study in *Solanum* was made was grown at Aroostook Farm of the MAINE AGRICULTURAL EXPERIMENT STATION by Doctor CHARLES F. CLARK of the Office of Horticultural Investigations, UNITED STATES DEPARTMENT OF AGRICULTURE. Doctor CLARK not only very kindly allowed the writer to have access to the material but also checked over the classification of the various forms and furnished information regarding the original sources of the wild species referred to in this paper. *S. chacoense* Bitter originally came from Paraguay, *S. demissum* Lindl. from Mexico, and *S. Fendleri* Gray and

S. Jamesii Torr. from the Alamo National Forest at Cloudcroft, New Mexico. The material was fixed in ALLEN's modification No. 15 of BOUIN's killing fluid. The cytological problem in this paper was suggested to the writer by Doctor KARL SAX of the MAINE AGRICULTURAL EXPERIMENT STATION. Doctor SAX also very kindly assisted the writer during the course of the study.




FIGURE 1.—Haploid chromosome group of *Solanum Jamesii* at heterotypic metaphase.




FIGURE 2.—Haploid chromosome group of *Solanum chacoense* at heterotypic metaphase.

As far as is now known¹ the chromosome number of all the cultivated varieties of *Solanum tuberosum* L. and the allied wild species is some multiple of 12. Figures 1 and 2 illustrate that the reduced number of




FIGURE 3.—Haploid Chromosome group of *Solanum Fendleri* at heterotypic metaphase.




FIGURE 4.—Haploid chromosome group of *Solanum demissum* at heterotypic metaphase.

chromosomes in *Solanum Jamesii* and *Solanum chacoense* respectively is 12. The chromosomes at homotypic metaphase in haploid cells of *Solanum Jamesii* are shown also in plate 1, figure 1. For *Solanum Fendleri*, as shown in figure 3 and plate 1, figure 2, the haploid chromosome number is 24, and for *Solanum demissum* as illustrated in figure 4 and plate 1, figure 3, the haploid chromosome number is 36.

In the cultivated varieties McIntyre and McCormick, 24 is the reduced number of chromosomes. Figure 5 and plate 1, figures 4 and 5 illustrate the heterotypic metaphase as observed in pollen mother cells of the McIntyre variety. There can be no question about the count in this variety as both poles clearly show 24 chromosomes. The two equatorial



FIGURE 5.—Haploid chromosome groups of the McIntyre variety at homotypic metaphase.

plates are approximately in one plane within the cell. The same clear-cut evidence has been obtained for the McCormick variety, for which figures 6 and 7 represent the heterotypic metaphase. In this case the chromosomes at the two poles were necessarily shown in separate drawings because the two plates were in different planes, one directly beneath the other.




FIGURE 6.—Haploid chromosome group of the McCormick Variety at homotypic metaphase.

FIGURE 7.—Haploid chromosome group of the McCormick variety at homotypic metaphase.

It is believed that tetraploidy occurs in the Early Ohio variety. Figures 8 and 9 and plate 1, figures 6 and 7 present evidence for this belief. These figures illustrate a pollen mother cell of the Early Ohio variety at heterotypic metaphase. Since both poles of the division figure show unquestionably more than 36 and approximately 48 chromosomes, there can be little doubt that these figures indicate tetraploidy in this variety.

LEGEND FOR PLATE 1

FIGURE 1.—Photomicrograph showing the haploid chromosome group of *Solanum Jamesii* at heterotypic metaphase. The 12 chromosomes of this species are clearly illustrated.

FIGURE 2.—Photomicrograph showing the haploid chromosome group of *Solanum Fendleri* at heterotypic metaphase. The 24 chromosomes of this species are clearly illustrated.


FIGURE 3.—Photomicrograph showing the haploid chromosome group of *Solanum demissum* at heterotypic metaphase. The 36 chromosomes of this species are clearly illustrated.

FIGURES 4 AND 5.—Photomicrographs of a pollen mother cell of the McIntyre variety of the cultivated potato at homotypic metaphase; in figure 4 one of the two equatorial plates is in focus and in figure 5 the other equatorial plate is in focus. These figures show that the haploid chromosome number of this variety is 24, and that the diploid chromosome number is 48.

FIGURES 6 AND 7.—Photomicrographs showing tetraploidy in the Early Ohio variety; figure 6 shows one of the two poles of the homotypic metaphase, and figure 7 shows the other pole of the same cell in focus.

FIGURE 8.—Photomicrograph showing what is believed to represent tetraploidy in the Early Rose variety. This figure shows the 48 chromosomes at heterotypic metaphase of the pollen mother cells.

SMITH, HUGH B. CHROMOSOME COUNTS IN *Solanum tuberosum*. PLATE 1.





Digitized by the Internet Archive
in 2026

https://archive.org/details/IA41551510_0106

At any rate the chromosome number is greater than a triploid form would have.




FIGURE 8.—Haploid chromosome group in the Early Ohio variety at homotypic metaphase, indicating tetraploidy in this variety.



FIGURE 9.—Haploid chromosome group in the Early Ohio variety at homotypic metaphase, indicating tetraploidy in this variety.

Although the evidence is not so conclusive as that which has just been described for the McIntyre and McCormick varieties, it is, nevertheless, believed that the haploid chromosome number in the Russet Rural and Early Ohio varieties is 24, and that tetraploidy, similar to that in Early Ohio, occurs in the Early Rose and Russet Rural varieties. The evidence




FIGURE 10.—Haploid chromosome group in the Russet Rural variety at heterotypic metaphase, indicating tetraploidy in this variety.



FIGURE 11.—Haploid chromosome group in the Russet Rural Variety at heterotypic metaphase.

for these tentative conclusions is presented in the following illustrations of heterotypic metaphases. Figures 10 and 11 illustrate respectively the haploid chromosome number of tetraploid and diploid plants of the Russet Rural variety. Figure 12 illustrates the haploid chromosome

number of diploid Early Ohio plants, and figures 13 and plate 1, figure 8 illustrate a pollen mother cell of the Early Rose variety at homotypic metaphase and are believed to indicate tetraploidy in this variety.

LUTMAN (1925) reports for the Irish Cobbler, Green Mountain, Lookout Mountain, and Early Rose varieties that the diploid chromosome number, as determined from a study of root tip material, is 36. This does not mean that heterotypic figures would necessarily show less than 24 chromosomes, as 12 may be single chromosomes and 12 of them paired. If this were the case, one would never be able to find 24 chromosomes at both poles of the heterotypic metaphase in these four varieties, and no such counts have been reported.




FIGURE 12.—Haploid chromosome group in the Early Ohio variety at heterotypic metaphase.




FIGURE 13.—Haploid chromosome group in the Early Rose variety at heterotypic metaphase, indicating tetraploidy in this variety.

If certain potato varieties are triploid, as is suggested by a count of 36 in somatic cells, one might find in this fact an explanation of at least part of the pollen sterility in the potato. Such a cause of sterility has been shown for *Oenothera* by GATES (1915), for *Datura* by BLAKESLEE, BELLING, and FARNUM (1920), for wheat by SAX (1921, 1922, 1922), and for roses by BLACKBURN AND HARRISON (1921). BLAKESLEE, BELLING, and FARNUM have shown that *Datura* mutants have a relatively high degree of pollen sterility and that the tetraploid form of *Datura Stramonium* is sterile with its parent stock. GATES states that only 11 percent of the pollen grains are viable in *Oenothera gigas* $\times$ *lata rubricalyr*. The irregular, 5- and 6-lobed grains, together with many other misshapen and sterile grains, may contain chromosome combinations which are incompatible with development. BLACKBURN AND HARRISON in their work with roses have found that univalents split at heterotypic metaphase and in many cases fail to reach the poles but form supernumerary micronuclei. Thus octads rather than tetrads are often formed and those pollen grains derived chiefly from micronuclei collapse. Such behavior is believed to

be due to hybridity. SAX working with wheat has found that the homologous chromosomes of the bivalents in species hybrids separate and pass regularly to the two poles, and univalents, if present, are distributed irregularly to one pole or the other during meiosis. He accounts for sterility in such hybrids on a hypothesis involving, "(1) the numerical and unbalanced relations of the chromosomes resulting from irregular meiotic divisions and (2) the specific interrelations of the parental chromosomes." He believes that the gametes exhibit more and more perfection as their chromosome constitution approaches more and more closely to that of the parental forms.

It has been shown by STOUT and CLARK (1924) that certain of the cultivated potato varieties have more pollen sterility than others. Perhaps the most sterile varieties have 12 univalent chromosomes and have a somatic chromosome number 36 and the least sterile ones have only paired chromosomes and have a diploid chromosome number of 48. The McIntyre and McCormick varieties do not have unpaired chromosomes and the haploid chromosome number in these two varieties is unquestionably 24; therefore, their 2X chromosome number is 48. It is of interest to note that STOUT and CLARK (1924) classify these two varieties as having a relatively high percentage of fertile pollen. Work is in progress to determine whether or not any of the commercial varieties exhibit lagging chromosomes during meiosis.

In this preliminary paper it has been shown that the haploid chromosome number is 12 for *S. Jamesii* and *S. chacoense*, 24 for *S. Fendleri*, and 36 for *S. demissum*. This is evidence that in these wild species of potato the haploid chromosome number is some multiple of 12, just as has been reported for other species of the Solanaceae. The McIntyre and McCormick cultivated varieties, which are characterized by a relatively high percentage of fertile pollen, have a haploid chromosome number of 24 and do not have unpaired chromosomes. Tetraploidy has occurred in the Early Ohio variety as shown by the appearance of haploid cells with approximately 48 chromosomes. The fact presented in this paper that the haploid chromosome number of potato has increased from 12 to 48 indicates that tetraploidy may have been a factor in the development of the cultivated varieties.

LITERATURE CITED

- BLACKBURN, KATHLEEN B., and HARRISON, J. W. H., 1921 The status of the British rose forms as determined by their cytological behavior. *Annals of Bot.* 35: 160-188.
BLAKESLEE, A. F., 1921 Types of mutation and their possible significance in evolution. *Amer. Nat.* 55: 254-267.

- BLAKESLEE, A. F., and BELLING, JOHN, 1924 Chromosomal chimeras in the Jimson weed. Science N.S. **60**: 19-20.
- BLAKESLEE, A. F., BELLING, JOHN, and FARNUM, M. E., 1920 Chromosomal duplication and Mendelian phenomena in *Datura* mutants. Science N.S. **52**: 388-390.
- BÖNICKE, L. V., 1911 Zur Kenntnis der Prophasen der heterotypischen Teilung einiger Pollenmutterzellen. Bericht. Deutsch. Bot. Gesell. **29**: 59-65.
- GATES, R. R., 1915 The mutation factor in evolution. London, Macmillan and Co.
- NĚMEC, B., 1899 Ueber Kern-und Zelltheilung bei *Solanum tuberosum*. Flora **86**: 214-227.
- SAX, KARL, 1921 Sterility in wheat hybrids. I. Sterility relationships and endosperm development. Genetics **6**: 229-416.
- 1922 Sterility in wheat hybrids. II. Chromosome behavior in partially sterile hybrids. Genetics **7**: 513-552.
- 1922 Sterility in wheat hybrids. III. Endosperm development and F_2 sterility. Genetics **7**: 553-558.
- STOUT, A. B., and CLARK, C. F., 1924 Sterilities of wild and cultivated potatoes with reference to breeding from seed. U. S. Dept. Agric. Bull. 1195: 1-32.
- WINKLER, H., 1910 Über die Nachkommenschaft der *Solanum* Pfropfbastarde und die Chromosomenzahlen ihrer Keimzellen. Zeitschr. f. Bot. **2**: 1-38.
- 1916 Über die experimentalle Erzeugung von Pflanzen mit abweichenden Chromosomenzahl. Zeitschr. f. Bot. **8**: 417-531.

INFORMATION FOR CONTRIBUTORS

The length of contributions to GENETICS will not be arbitrarily limited but articles of less than fifty pages may be expected to appear more promptly than those of greater length. The limitations as to subject matter will be construed broadly, so that any paper of sufficient importance, whether in the field of genetics proper, of cytology, embryology, physiology, biometry, or mathematics, if of such character as to be of primary interest to the geneticist, will find here an appropriate place of publication.

Contributors are requested to observe the usual care in the preparation of manuscripts. All references to literature should cite the name of the author, followed by the year of publication, the papers so referred to being collected into a list of "Literature cited" at the end of the article. In this list great care should be taken to give titles *in full*, and to indicate accurately in Arabic numerals, the volume number, and first and last pages, and the date of publication, of each paper, if published in a periodical, and the number of pages, place and date of publication, and name of publisher, of each independent publication. The arrangement of this list should be alphabetical by authors, and chronological under each author.

Foot-notes, legends for figures, tabular matter, and quotations involving more than one sentence should be typewritten on separate sheets. All foot-notes should be numbered consecutively in a single series and designated by Arabic superscript numerals.

No limitation will be placed on the kind of illustration to be used, but as funds for illustrations are still inadequate, authors are requested to practice restraint, and to limit themselves to such amount and grade of illustration as seem positively essential to an adequate understanding of their work. The size of the typeforms of GENETICS, exclusive of the headings, is $4\frac{5}{8} \times 7$ inches and all illustrations intended for use as single plates should not exceed these dimensions or should allow of reduction to them. Folded plates may be of any desired length, horizontally, but are limited to 7 inches vertically.

Authors will receive galley proofs, and, if specially requested, page proofs will be sent also, it being understood that the latter must be returned very promptly, and that no extensive change may be made in the page proofs, which is not compensated for within the same paragraph, or in an adjacent paragraph on the same page. Changes from copy will be charged against the author.

Contributors will receive 50 reprints of their papers, with covers, free of charge, provided not fewer than 100 are ordered. If a less number is desired, one-half the total cost of their manufacture will be charged to the author. Additional copies may be secured at actual cost of manufacture, provided these are ordered when the corrected galley proofs are returned.

Manuscripts and all editorial correspondence should be addressed to the Managing Editor, Dr. DONALD F. JONES, P. O. Drawer 1106, New Haven, Connecticut.

GENETICS, JANUARY 1927

TABLE OF CONTENTS

	PAGE
LI, JU-CHI, The effect of chromosone aberrations in <i>Drosophila melanogaster</i>	1
SOROKIN, HELEN, Cytological and morphological investigations on gynodimorphic and normal forms of <i>Ranunculus acris</i> L.....	59
SMITH, HUGH B., Chromosome counts in the varieties of <i>Solanum tuberosum</i> and allied wild species.....	84

FRONTISPIECE

In the death of WILLIAM BATESON on February 8, 1926, the science of genetics has suffered its greatest loss since the Mendelian renaissance in 1900. The Editorial Board of GENETICS takes this occasion to express its profound sense of appreciation of the splendid contributions both of fact and of spirit which BATESON made to the cause of genetical discovery. The Editorial Board had voted in 1921 to publish a portrait of BATESON as frontispiece to volume 8 of GENETICS, but as no satisfactory portrait could be found at that time the matter was postponed. The Board has thus the satisfaction of having offered its tribute to the living.

The photograph here reproduced was taken at Christmas time, 1922, and is pronounced by Mrs. BATESON to be, in her judgment, the best available portrait of her husband. We are deeply indebted to Mrs. BATESON for copies of this photograph and also for biographical data to be used in preparing a sketch of BATESON's life which will accompany the title pages of this volume of GENETICS.

The cost of engraving and printing this portrait has been generously contributed by Doctor JOHN HARVEY KELLOGG, well-known Superintendent of the BATTLE CREEK SANITARIUM, Battle Creek, Michigan, whose interest in eugenics led him to establish, about twenty years ago, a corporation known as the RACE BETTERMENT FOUNDATION, which organized the first RACE BETTERMENT CONGRESS in 1914, and which established in 1915 an "EUGENICS REGISTRY." The RACE BETTERMENT FOUNDATION of which Doctor KELLOGG is President has recently established BATTLE CREEK COLLEGE, one of whose major aims is the promotion of eugenics and race betterment ideals.

[This number of GENETICS (12 : 1-92) was issued May 20, 1927]